



Use Lamisil DermGel with caution:

- If your skin is inflamed or broken.
Lamisil DermGel contains alcohol and may be irritating to damaged skin.

Consult your doctor or pharmacist before using Lamisil DermGel:

- If you are pregnant, think you might be pregnant or are trying to become pregnant.
Lamisil DermGel should not be used during pregnancy unless your doctor has advised you to.

Taking other medicines

Tell your doctor or pharmacist if you are using any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may affect the way other medicines work. Do not apply other medicinal products to the areas being treated.

Important information about some of the ingredients of Lamisil DermGel

Lamisil DermGel contains butylhydroxytoluene, which may cause local skin reactions (e.g. contact dermatitis), or irritation to the eyes and mucous membranes.

USING LAMISIL DERMGEL

How to use it

1. Clean and dry the affected area.
2. Unscrew the cap, and if using for the first time, use the point in the cap to pierce the top of the tube, then gently squeeze the tube.
3. Apply a thin layer of Lamisil DermGel to the affected skin and surrounding area.
4. Rub in gently.
5. Replace the cap on the tube.

6. Wash your hands after touching the infected skin so that you do not spread the infection.

Lamisil DermGel is for external use only. Do not use in the mouth or swallow it. Avoid applying the gel near your eyes or touching your eyes while you still have gel on your hands. In case of accidental contact with the eyes, rinse the eyes thoroughly with running water and if any symptoms persist see your doctor.

If you are treating an infection in a skin-fold area such as under the breast, armpit or groin region, you can cover the treated area with gauze. This is particularly useful at night. If you do, use clean gauze each time you apply the gel.

How long to use it for

Apply Lamisil DermGel to the infected areas once per day for 7 days.

It is important to use Lamisil DermGel for the recommended treatment period, even if the symptoms disappear earlier. Infections usually appear to improve within a few days, but may return if the gel is not applied as directed.

If you have not noticed any signs of improvement within 2 weeks of first starting treatment, see your doctor or pharmacist.

If you forget to use it

If you forget an application of Lamisil DermGel, apply it as soon as you remember and then go back to using it as you would normally. If it is almost time for your next application, skip the application you missed and apply the usual amount of Lamisil DermGel when you are next meant to. Do not use a double dose to make up for the dose that you missed.

2003847_1 / 5055610 AU/NZ



WHAT LAMISIL DERMGEL IS USED FOR

Lamisil DermGel is used to treat fungal infections of the skin including the following tinea infections:

- Athlete's foot - between the toes
- Jock itch - groin and inner thigh area
- Ringworm - commonly affects arms, legs, trunk or neck

Lamisil DermGel can also be used to treat skin infections caused by Pityriasis versicolor, a superficial fungal infection of the skin.

You should apply Lamisil DermGel only to the skin on the outside of the body. This product is not suitable for treating fungal infections of the nail.

BEFORE USING LAMISIL DERMGEL

Do not use Lamisil DermGel:

- If you are allergic to terbinafine or to any other ingredient listed at the end of this leaflet
- If you are breastfeeding
- On children younger than 18 years or allow them to come into contact with treated skin.



Try to use Lamisil DermGel as directed. This is important as missing applications may cause the infection to return.

SIDE EFFECTS

Tell your doctor or pharmacist if you do not feel well while you are using Lamisil DermGel.

Most people benefit from using Lamisil DermGel but it may have unwanted side effects in a few people.

Ask your doctor or pharmacist to answer any questions you may have.

Stop using Lamisil DermGel immediately and talk to your doctor if you have any of the following which may be signs of an allergic reaction:

- difficulty in breathing or swallowing
- swelling of the face, lips, tongue or throat
- severe itching of the skin, with a red rash or raised bumps

Some side effects are common. These include:

- skin peeling
- itching

Some side effects are uncommon. These include:

- skin lesion
- scab
- skin disorder
- skin colour changes
- redness
- burning sensation
- pain
- application site pain or irritation

Some side effects are rare. These include:

- dry skin
- eczema

If Lamisil DermGel is accidentally applied to the eyes, eye irritation may occur.

If you get any other side effects, talk to your doctor or pharmacist. This may include side effects not listed in this leaflet.

IN CASE OF OVERDOSE

If Lamisil DermGel is accidentally swallowed, particularly by a child, contact a doctor or the Poisons Information Centre by calling 13 11 26 in Australia or 0800 764 766 in New Zealand.

AFTER USING LAMISIL DERMGEL

Storage

- Store below 25°C.
- Store out of reach of children.
- Do not use past the expiry date shown on the pack.
- Store in the original package.

Disposal

If your pharmacist or doctor tells you to stop using Lamisil DermGel or you find that it has passed the expiry date, ask your pharmacist what to do with any that you have left over.

PRODUCT DESCRIPTION

What it looks like

Lamisil DermGel is a white gel, which is available in a 15 g varnished aluminium or laminate tube.

Lamisil DermGel is enclosed in tamper-evident packaging to help protect you from product tampering. If you find that the tube seal has been broken or is missing, do not use the product and return it to your pharmacist.



Ingredients

Lamisil DermGel contains the active ingredient terbinafine hydrochloride.

It also contains the following inactive ingredients:

- purified water
- ethanol
- isopropyl myristate
- polysorbate 20
- carbomer 934P
- sorbitan monolaurate
- benzyl alcohol
- sodium hydroxide
- butylated hydroxytoluene

Distributed by:

Nice Pak Products Pty Ltd
120 Woodlands Drive
Braeside, VIC 3195, Australia.

Nice Pak Products (NZ) Ltd
PO Box 37217
Parnell, Auckland 1151, New Zealand.



AU: 1800 506 750
NZ : 0800 555 895

This leaflet was last revised in March 2024